

Explicit expressions for meromorphic solution of autonomous nonlinear ordinary differential equations

Maria V. Demina, Nikolay A. Kudryashov

Department of Applied Mathematics, National Research
Nuclear University MEPhI, 31 Kashirskoe Shosse, 115409
Moscow, Russian Federation

Abstract

Meromorphic solutions of autonomous nonlinear ordinary differential equations are studied. An algorithm for constructing meromorphic solutions in explicit form is presented. General expressions for meromorphic solutions (including rational, periodic, elliptic) are found for a wide class of autonomous nonlinear ordinary differential equations.

1 Introduction

Autonomous nonlinear partial and ordinary differential equations frequently arise in mathematical models describing processes and phenomena in physics, chemistry, biology, etc. The problem of constructing exact solutions to autonomous nonlinear differential equations is intensively studied [1–10]. A number of methods and algorithms has been introduced. As a rule these methods deal with a priori fixed families of meromorphic exact solutions. Consequently, solutions outside given families may be missed. Very often various methods yield the same families of exact solutions written in a different way. And as a result many "new" exact solutions appear. [11–14] Thus an important problem is to classify families of exact solutions. In this paper we study the general form of meromorphic solutions to autonomous nonlinear ordinary differential equations. The main aim is to present our algorithm, which can be applied to construct exact meromorphic solutions in explicit form. For a wide class of nonlinear ordinary differential equations we make

a conclusion that exact solutions given in our paper are the only possible meromorphic solutions.

This paper is organized as follows. In section 2 we describe our method and present explicit expressions for meromorphic solutions. In sections 3 we give an example and construct meromorphic solutions of a certain second order nonlinear differential equation.

2 Method applied

Let us look for nonconstant meromorphic exact solutions of an autonomous nonlinear ordinary differential equation

$$E[w(z)] = 0. \quad (1)$$

In this expression $E[w(z)]$ is a polynomial in $w(z)$ and its derivatives. For every solution $w(z)$ of equation (1) there exists a family of solutions $w(z - z_0)$. Without loss of generality, we omit arbitrary constant z_0 . Suppose the following.

Condition I. For solutions of equation (1) there are only N different asymptotic expansions corresponding to the Laurent series in a neighborhood of $z = 0$

$$w^{(i)}(z) = \sum_{k=1}^{p_i} \frac{c_{-k}^{(i)}}{z^k} + \sum_{k=0}^{\infty} c_k^{(i)} z^k, \quad 0 < |z| < \varepsilon_i, \quad i = 1, \dots, N. \quad (2)$$

In this expression $p_i > 0$ is an order of the pole $z = 0$. If $N = 1$ we shall omit the upper index.

Condition II. Substituting $w(z) = \lambda W(z)$ into equation (1) yields expression with only one term of the highest degree in respect of λ .

Note that if $w(z)$ has more than N poles, then $w(z)$ is periodic. Indeed, there exist poles $z = z_1, z = z_2$ of $w(z)$ such that the functions $w(z + z_1), w(z + z_2)$ are meromorphic solutions of equation (1) with a pole at $z = 0$. Thus we have the equality $w(z) = w(z + z_2 - z_1)$.

Meromorphic solutions of equation (1) are classified in the following theorems.

Theorem 1. All meromorphic solutions of equation (1) satisfying condition I with $N = 1$ and condition II are of the form:

1) elliptic solutions with the periods $2\omega_1, 2\omega_2$

$$w(z) = \left\{ \sum_{k=2}^p \frac{(-1)^k c_{-k}}{(k-1)!} \frac{d^{k-2}}{dz^{k-2}} \right\} \wp(z; \omega_1, \omega_2) + h_0, \quad (3)$$

Necessary condition for elliptic solutions to exist is $c_{-1} = 0$.

2) periodic solutions with the period T

$$w(z) = \frac{\pi}{T} \left\{ \sum_{k=1}^p \frac{(-1)^{k-1} c_{-k}}{(k-1)!} \frac{d^{k-1}}{dz^{k-1}} \right\} \cot\left(\frac{\pi z}{T}\right) + h_0. \quad (4)$$

3) rational solutions

$$w(z) = \sum_{k=1}^p \frac{c_{-k}}{z^k} + \sum_{k=0}^m c_k z^k, \quad m \geq 0. \quad (5)$$

In 1) and 2) h_0 is a constant.

Proof. For any meromorphic function $f(z)$ with the poles $\{a_n\}$ Mittag-Leffler found the representation [15]

$$f(z) = \sum_{n=1}^{\infty} \left\{ P_n \left(\frac{1}{z - a_n} \right) - g_n(z) \right\} + h(z), \quad (6)$$

where $\{P_n[(z - a_n)^{-1}]\}$ are principal parts of Laurent expansions for the function $f(z)$ around poles $\{a_n\}$, $\{g_n(z)\}$ are polynomials, and $h(z)$ is an entire function. The polynomial $g_n(z)$ is added to the n^{th} term of the sum (6) in order to provide convergence of the series. A series of the form (6) is said to be convergent in $M \subset \mathbb{C}$ if only finite number of terms possesses poles in M and without these terms the series converges [15]. Suppose $w(z)$ has two or more poles. Then from condition I follows that $w(z)$ is doubly periodic (elliptic) or simply periodic. For the elliptic function $w(z)$ with Laurent expansion (2) expression (6) reads

$$w(z) = \frac{c_{-2}}{z^2} + \sum_{(n,m) \neq (0,0)} \left[\frac{c_{-2}}{(z - 2n\omega_1 - 2m\omega_2)^2} - \frac{c_{-2}}{(n\omega_1 + m\omega_2)^2} \right] + \sum_{k=3}^p \sum_{n,m} \frac{c_{-k}}{(z - 2n\omega_1 - 2m\omega_2)^k} + h_0. \quad (7)$$

Solution (7) can be rewritten in terms of Weierstrass elliptic function $\wp$ satisfying the equation

$$(\wp_z)^2 = 4\wp^3 - g_2\wp - g_3. \quad (8)$$

Thus we obtain representation for the elliptic solution $w(z)$ in the form (3). Necessary condition $c_{-1} = 0$ follows from the theorem for total sum of the

residues of an elliptic function in the parallelogram of periods. Writing expression (6) for the solution $w(z)$ with period T yields

$$w(z) = \frac{c_{-1}}{z} + \sum_{n \neq 0} \left[\frac{c_{-1}}{(z - nT)} + \frac{c_{-1}}{nT} \right] + \sum_{k=2}^p \sum_n \frac{c_{-k}}{(z - nT)^k} + h(z). \quad (9)$$

In the paper [16] meromorphic traveling wave solutions of the Kuramoto – Sivashinsky equation were studied. A powerful method based on Nevanlinna theory [17] was developed. From the results of the paper [16] it follows that periodic solutions of equation (1) satisfying condition II are rational in e^{rz} . Making change of variables $s = e^{rz}$, $y(s) = w(z)$ in equation (1) and then substituting $y = s^n$, $n > 0$ as s tends to infinity and $y = s^{-l}$, $l > 0$ as s tends to zero into the result, we see that an entire periodic function $h(z)$ in (9) is a constant since condition II is valid. Further, making use of the formula

$$\frac{1}{z} + \sum_{n \neq 0} \left[\frac{1}{(z - nT)} + \frac{1}{nT} \right] = \frac{\pi}{T} \cot \left(\frac{\pi z}{T} \right) \quad (10)$$

we get (4) from (9) with $h(z) = h_0$. Now suppose $w(z)$ has only one pole, then

$$w(z) = \sum_{k=1}^p \frac{c_{-k}}{z^k} + h(z), \quad (11)$$

where an entire function $h(z)$ does not have an essential singularity at infinity [16]. This completes the proof.

Remark. For fixed values of parameters in equation (1), if any, there may exist only one meromorphic solution (rational, simply periodic or elliptic) with a pole at $z = 0$.

In the next theorem we generalize results of theorem 1 to the case $N > 1$. Let us call poles of meromorphic solution $w(z)$ with the Laurent expansion $w^{(i)}(z)$ (see (2)) as poles of type i .

Theorem 2. All meromorphic solutions of equation (1) satisfying condition I and condition II are of the form:

1) elliptic solutions with the periods $2\omega_1, 2\omega_2$

$$w(z) = \left\{ \sum_{i \in I} \sum_{k=2}^{p_i} \frac{(-1)^k c_{-k}^{(i)}}{(k-1)!} \frac{d^{k-2}}{dz^{k-2}} \right\} \left(\frac{1}{4} \left[\frac{\wp_z(z) + B_i}{\wp(z) - A_i} \right]^2 - \wp(z) \right) + \sum_{i \in I} \frac{c_{-1}^{(i)} (\wp_z(z) + B_i)}{2 (\wp(z) - A_i)} + \left\{ \sum_{k=2}^{p_{i_0}} \frac{(-1)^k c_{-k}^{(i_0)}}{(k-1)!} \frac{d^{k-2}}{dz^{k-2}} \right\} \wp(z) + h_0, \quad (12)$$

where $\wp(z) \stackrel{def}{=} \wp(z; \omega_1, \omega_2)$, $B_i^2 = 4A_i^3 - g_2A_i - g_3$. Necessary condition for elliptic solutions (12) to exist is

$$\sum_{i \in I} c_{-1}^{(i)} + c_{-1}^{(i_0)} = 0. \quad (13)$$

2) periodic solutions with the period T

$$\begin{aligned} w(z) = & \frac{\pi}{T} \left\{ \sum_{i \in I} \sum_{k=1}^{p_i} \frac{(-1)^{k-1} c_{-k}^{(i)}}{(k-1)!} \frac{d^{k-1}}{dz^{k-1}} \right\} \frac{A_i \cot\left(\frac{\pi z}{T}\right) + \frac{\pi}{T}}{A_i - \frac{\pi}{T} \cot\left(\frac{\pi z}{T}\right)} \\ & + \frac{\pi}{T} \left\{ \sum_{k=1}^{p_{i_0}} \frac{(-1)^{k-1} c_{-k}^{(i_0)}}{(k-1)!} \frac{d^{k-1}}{dz^{k-1}} \right\} \cot\left(\frac{\pi z}{T}\right) + h_0. \end{aligned} \quad (14)$$

3) rational solutions

$$w(z) = \sum_{k=1}^{p_{i_0}} \frac{c_{-k}^{(i_0)}}{z^k} + \sum_{i \in I} \sum_{k=1}^{p_i} \frac{c_{-k}^{(i)}}{(z - a_i)^k} + \sum_{k=0}^m h_k z^k, \quad m \geq 0. \quad (15)$$

In 1), 2) and 3) $\{A_i\}$, $\{B_i\}$, $\{a_i\}$, $\{h_i\}$ are constants and $I = \emptyset$ or $I \subseteq \{1, 2, \dots, N\} \setminus \{i_0\}$, $1 \leq i_0 \leq N$.

Proof. Let $w(z)$ be a meromorphic solution of equation (1) with poles of types j , $j \in J$, $J \subseteq \{1, 2, \dots, N\}$. Without loss of generality, we may select any $i_0 \in J$ and suppose that the point $z = 0$ is a pole of type i_0 for $w(z)$. Let us set $I = J \setminus \{i_0\}$. For $N = 1$ we see that $I = \emptyset$. Let us consider the case $N > 1$. If there are more than $|I| + 1$ poles of different types, then $w(z)$ is periodic (simply periodic or elliptic). By analogy with the case $N = 1$ we get the expression for elliptic solutions

$$w(z) = \sum_{i \in I \cup \{i_0\}} c_{-1}^{(i)} \zeta(z - a_i) + \left\{ \sum_{i \in I \cup \{i_0\}} \sum_{k=2}^{p_i} \frac{(-1)^k c_{-k}^{(i)}}{(k-1)!} \frac{d^{k-2}}{dz^{k-2}} \right\} \wp(z - a_i) + \tilde{h}_0, \quad (16)$$

where $\zeta(z)$ is the Weierstrass ζ -function and a_{i_0} may be taken as zero. Note that $a_i \neq a_j$, $i, j \in I \cup \{i_0\}$, $i \neq j$ are poles of elliptic solutions (16) in the parallelogram of periods built on $2\omega_1$, $2\omega_2$. The function $\zeta(z)$ has a simple pole at the point $z = 0$ with residue 1. The theorem for total sum of the residues of an elliptic function in the parallelogram of periods yields condition

(13). Using the addition formulae [18]

$$\begin{aligned}\zeta(z - a_i) &= \zeta(z) - \zeta(a_i) + \frac{1}{2} \frac{\wp_z(z) + \wp_z(a_i)}{\wp(z) - \wp(a_i)}, \\ \wp(z - a_i) &= -\wp(z) - \wp(a_i) + \frac{1}{4} \left[\frac{\wp_z(z) + \wp_z(a_i)}{\wp(z) - \wp(a_i)} \right]^2\end{aligned}\tag{17}$$

and notation $A_i \stackrel{\text{def}}{=} \wp(a_i)$, $B_i \stackrel{\text{def}}{=} \wp_z(a_i)$, we get equality (12). Further, from equation (8) we have $B_i^2 = 4A_i^3 - g_2A_i - g_3$. Note that for the parameters h_0 and $\tilde{h}_0$ the following correlation holds

$$h_0 = \tilde{h}_0 - \sum_{i \in I} c_{-1}^{(i)} \zeta(a_i) - \sum_{i \in I} c_{-2}^{(i)} \wp(a_i).\tag{18}$$

Similarly to the case $N = 1$ simply periodic solutions can be written as

$$w(z) = \frac{\pi}{T} \left\{ \sum_{i \in I \cup \{i_0\}} \sum_{k=1}^{p_i} \frac{(-1)^{k-1} c_{-k}^{(i)}}{(k-1)!} \frac{d^{k-1}}{dz^{k-1}} \right\} \cot \left(\frac{\pi(z - a_i)}{T} \right) + h_0,\tag{19}$$

where again we may set $a_{i_0} = 0$ and $a_i \neq a_j$, $i, j \in I \cup \{i_0\}$, $i \neq j$ are poles of simply periodic solutions (16) in a stripe built on T . With the help of addition formulae for $\cot[\pi(z - a_i)T^{-1}]$ we obtain expression (14), where $A_i \stackrel{\text{def}}{=} \pi \cot[\pi a_i T^{-1}] T^{-1}$. For rational solutions we have the equality

$$w(z) = \sum_{i \in I \cup \{i_0\}} \sum_{k=1}^{p_i} \frac{c_{-k}^{(i)}}{(z - a_i)^k} + \sum_{k=0}^m h_k z^k, \quad m \geq 0,\tag{20}$$

where $a_i \neq a_j$, $i, j \in I \cup \{i_0\}$, $i \neq j$. Without loss of generality, we set $a_{i_0} = 0$ in (20) to obtain (15).

Remark 1. Again for fixed values of parameters in equation (1), if any, there may exist only one meromorphic solution (rational, simply periodic or elliptic) with a pole at $z = 0$ of type i , $1 \leq i \leq N$.

Remark 2. While making classification of meromorphic solutions, we should consider all the variants for $J \subseteq \{1, 2, \dots, N\}$.

Now let us discuss the problem of finding exact meromorphic solutions in explicit form. The main idea is to expand exact solutions given in theorems 1 and 2 around their poles and to compare coefficients of these expansions with coefficients of series (2). Our algorithm is the following.

At the *first step* one constructs asymptotic expansions corresponding to Laurent series in a neighborhood of a pole $z = 0$.

At the *second step* one selects an expression for a meromorphic solution (see theorem 1).

At the *third step* one expands the meromorphic solution in a neighborhood of poles $z = 0$, $z = a_i$, $i \in I$.

At the *fourth step* one compares coefficients of the series found at the first and the third steps and solves an algebraic system for the parameters of exact meromorphic solution. Along with this, correlations for the parameters of equation (1) may arise. If the meromorphic solution possesses poles of different types, it is convenient to consider all possible Laurent expansions. Optimal number of equations in algebraic system equals to the number of parameters of the meromorphic solution and equation (1) plus 1. If this system is inconsistent, then equation (1) does not possess meromorphic solutions with supposed expression.

At the *fifth step* one verifies the meromorphic solution obtained at the previous step, substituting this solution and correlations on the parameters of equation (1) into the latter.

In order to find the Laurent expansions for elliptic solutions (theorem 1) in a neighborhood of poles $z = 0$, $z = a_i$, $i \in I$ one takes formula (16) with $a_{i_0} = 0$ and then introduces notation $A_i \stackrel{\text{def}}{=} \wp(a_i)$, $B_i \stackrel{\text{def}}{=} \wp_z(a_i)$, $i \in I$. The parameters of elliptic solutions (12) that one needs to calculate are g_2 , g_3 , h_0 , A_i , $i \in I$.

As far as simply periodic solutions (theorem 1) are concerned, one takes formula (19) with $a_{i_0} = 0$, constructs Laurent expansions in a neighborhood of poles $z = 0$, $z = a_i$, $i \in I$, and introduces notation $L \stackrel{\text{def}}{=} \pi^2 T^{-2}$, $A_i \stackrel{\text{def}}{=} \pi \cot[\pi a_i T^{-1}] T^{-1}$, $i \in I$. The parameters of simply periodic solutions (14) to be found are the following L , h_0 , A_i , $i \in I$.

For rational solutions (theorem 2) the parameters to be calculated are h_0 , a_i , $i \in I$. In order to find the highest exponent m (see (15)) one may construct the Laurent expansion in a neighborhood of infinity.

To conclude this section we would like to emphasize that our method is also applicable to algebraic autonomous nonlinear ordinary differential equations with arbitrary coefficients in asymptotic expansions corresponding to the Laurent series in a neighborhood of poles. Arbitrary coefficients should be added to the list of parameters. However, these equations may possess meromorphic solutions of more complicated structure.

3 Explicit construction of meromorphic solutions

As an example let us find exact meromorphic solutions of the following second order autonomous nonlinear differential equation

$$w_{zz} + ww_z + \sigma w^3 + \delta w_z + \beta w^2 + \alpha w + \gamma = 0, \quad \sigma \neq 0. \quad (21)$$

Without loss of generality, we may set $\delta = 0$, $\sigma = -6$, and study the equation

$$w_{zz} + ww_z - 6w^3 + \beta w^2 + \alpha w + \gamma = 0. \quad (22)$$

This equation admits two different asymptotic expansions corresponding to Laurent series in a neighborhood of poles. They are the following

$$w^{(1)}(z) = \frac{1}{2z} + \frac{\beta}{20} + \left(\frac{11\beta^2}{200} + \alpha \right) \frac{z}{9} + \left(\frac{\beta^3}{375} + \frac{\alpha\beta}{15} + \gamma \right) \frac{z^2}{2} + o(|z|^2) \quad (23)$$

$$w^{(2)}(z) = -\frac{2}{3z} + \frac{2\beta}{33} - \left(\frac{20\beta^2}{363} + \alpha \right) \frac{z}{12} + \frac{3}{20} \left(\frac{2\beta^3}{1331} + 22\alpha\beta + \gamma \right) z^2 + o(|z|^2) \quad (24)$$

We see that equation (22) satisfies conditions II and I with $N = 2$. First let us construct simply periodic solutions possessing poles of single type (with residue $c_{-1}^{(1)} = 1/2$ and then with residue $c_{-1}^{(2)} = -2/3$). In this case we should take exact meromorphic solution in the form

$$w(z) = c_{-1}\sqrt{L} \cot\left(\sqrt{L}z\right) + h_0, \quad L \stackrel{\text{def}}{=} \frac{\pi^2}{T^2}. \quad (25)$$

Note that we omit arbitrary constant z_0 . Finding the Laurent expansion of function (25) at the point $z = 0$

$$w(z) = \frac{c_{-1}}{z} + h_0 - \frac{c_{-1}L}{3}z - \frac{c_{-1}L^2}{45}z^3 - \frac{2c_{-1}L^3}{945}z^5 + o(|z|^5) \quad (26)$$

and comparing coefficients of this series with coefficients of expansion (23) (and then (24)), we obtain parameters L , h_0 and correlations on the parameters α , β , γ . Substituting obtained meromorphic solutions into equation (22), we make sure that this equation indeed possesses meromorphic solutions of the form (25). For the Laurent series (23) these solutions can be written as

$$w(z) = \frac{\pi}{2T} \cot\left\{\frac{\pi(z-z_0)}{T}\right\} + \frac{\beta}{20}, \quad T = 10 \left(-\frac{3}{11\beta^2 + 200\alpha} \right)^{1/2} \pi \quad (27)$$




Figure 1: Family of solutions (27).

provided that the following correlation holds

$$\gamma = -\frac{(\beta^2 + 25\alpha)\beta}{375}. \quad (28)$$

For the Laurent series (24) we get the family of meromorphic solutions

$$w(z) = -\frac{2\pi}{3T} \cot \left\{ \frac{\pi(z - z_0)}{T} \right\} + \frac{2\beta}{33}, \quad T = 22 \left(-\frac{2}{20\beta^2 + 363\alpha} \right)^{1/2} \pi \quad (29)$$


and conditions on the parameters of equation (22)

$$\gamma = -\frac{(4\beta^2 + 121\alpha)\beta}{2662}. \quad (30)$$

Note that in expressions (27), (29) the period T depends on the parameters α and β and can be real or complex. For the functions with purely imaginary period $T = i\tau$, $\tau \in \mathbb{R}$ the following correlation holds

$$\frac{\pi}{T} \cot \left(\frac{\pi z}{T} \right) = \frac{\pi}{\tau} \coth \left(\frac{\pi z}{\tau} \right). \quad (31)$$

Now let us construct exact meromorphic solutions with poles of different types. The general expression for this solution is the following (see expression



(a) $\alpha = -10, \beta = 10, z_0 = 0, T = 22\pi/\sqrt{815}$ (b) $\alpha = 10, \beta = 10, z_0 = 0, T = 22\pi i/\sqrt{2815}$

Figure 2: Family of solutions (29).

(14) with $i_0 = 1, I = \{2\}, a_2 = a$)

$$w(z) = c_{-1}^{(1)}\sqrt{L}\cot\left\{\sqrt{L}z\right\} + c_{-1}^{(2)}\sqrt{L}\cot\left\{\sqrt{L}(z-a)\right\} + h_0. \quad (32)$$

Again we omit arbitrary constant z_0 . Without loss of generality, the point $z = 0$ is a pole of the first type, while the point $z = a$ is a pole of the second type. Constructing the Laurent series for solution (32) in a neighborhood of the points $z = 0, z = a$, we obtain

$$\begin{aligned} w(z) &= \frac{c_{-1}^{(1)}}{z} + h_0 - Ac_{-1}^{(2)} - \left(c_{-1}^{(2)}(L+A) + \frac{c_{-1}^{(1)}L}{3}\right)z + o(|z|), \\ w(z) &= \frac{c_{-1}^{(2)}}{z-a} + h_0 + Ac_{-1}^{(1)} - \left(c_{-1}^{(1)}(L+A) + \frac{c_{-1}^{(2)}L}{3}\right)(z-a) + o(|z-a|), \end{aligned} \quad (33)$$

where we set $A \stackrel{def}{=} \sqrt{L}\cot\sqrt{L}a$. Comparing the first of these series with the expansion (23) and the second — with expansion (24), we find the parameters h_0, T, A and conditions on the parameters α, β, γ of equation (22). Thus

we have

$$w(z) = \frac{\beta}{220} \cot \left\{ \frac{\beta(z - z_0)}{110} \right\} - \frac{\beta \left(7 \cot \left\{ \frac{\beta(z - z_0)}{110} \right\} + 1 \right)}{165 \left(\cot \left\{ \frac{\beta(z - z_0)}{110} \right\} + 7 \right)} + \frac{61\beta}{660}. \quad (34)$$

The family of meromorphic functions (34) solves equation (22) provided that

$$\alpha = -\frac{94\beta^2}{3025}, \quad \gamma = -\frac{31\beta^3}{33275}. \quad (35)$$

Expression (34) can be rewritten as

$$w(z) = \frac{\beta}{220} \cot \left\{ \frac{\beta(z - z_0)}{110} \right\} - \frac{\beta}{165} \cot \left\{ \frac{\beta(z - z_0)}{110} + \frac{1}{2} \log \frac{4}{3} \right\} + \frac{61\beta}{660}. \quad (36)$$

Meromorphic solutions (34) with $z_0 = 0$ possess two poles on real axes, one – at the origin, another one – at the point $a = -55 \log(4/3)/\beta$, $\beta \in \mathbb{R}$. All other poles lie in the complex domain since the period of solutions is $T = 110\pi i/\beta$.

Now let us construct rational solutions of the equation in question. Substituting $w = z^m$ as z tends to infinity into equation (22) we see that $m = 0$. Note that at certain conditions on the parameters α , β , γ expansions (23), (24) terminate. As a result we find two families of rational solutions and correlations for the parameters of equation (22)

$$\begin{aligned} w(z) &= \frac{1}{2(z - z_0)} + \frac{\beta}{20}, \quad \alpha = -\frac{11\beta^2}{200}, \quad \gamma = \frac{\beta^3}{1000} \\ w(z) &= -\frac{2}{3(z - z_0)} + \frac{2\beta}{33}, \quad \alpha = -\frac{20\beta^2}{363}, \quad \gamma = \frac{4\beta^3}{3993}. \end{aligned} \quad (37)$$

If we try to find rational solutions with poles of two types, we obtain an inconsistent system of algebraic equations. Consequently, rational solutions of the form

$$w(z) = \frac{c_{-1}^{(1)}}{z - z_0} + \frac{c_{-1}^{(2)}}{z - a - z_0} + h_0. \quad (38)$$

do not satisfy equation (22).

Equation (22) does not possess elliptic solutions, since necessary condition (13) does not hold.

We have studied all the expressions for meromorphic solutions given in theorem 2. Hence, we have found all families of nonconstant meromorphic solutions of equation (22). Plots of these solutions for certain values of the parameters are given in figures 1 — 3.




Figure 3: Family of solutions (34).

4 Conclusion

In this paper we have presented a method for constructing exact solutions of autonomous nonlinear differential equations. Our approach is simple in application since one can introduce parameters of exact solutions in such a way that for many equations linear correlations on the parameters arise at each step. Our method generalizes several other methods (for example, exp-function method, tanh-function method [19] and their extensions and modifications [20]). A great drawback of these methods is an a priori assumed expression for an exact solution. As a consequences solutions outside stated family may be lost. Our approach takes into account the structure of singularities and possible Laurent expansions for exact solutions. Along with this, we have obtained the general form of meromorphic solutions for a wide class of autonomous nonlinear ordinary differential equations.

5 Acknowledgements

This research was partially supported by Federal Target Programm "Research and Scientific - Pedagogical Personnel of innovation in Russian Federation on 2009 – 2013 (Contract P2457).

References

- [1] Kudryashov N.A. Exact soliton solutions of the generalized evolution equation of wave dynamics. J Appl Math and Mech 1988; 52(3): 360–5.
- [2] Kudryashov N.A. Exact solutions of the generalized Kuramoto–Sivashinsky equation, Phys Lett A 1990; 147: 287–91.
- [3] Kudryashov N.A. On types of nonlinear nonintegrable differential equations with exact solutions, Phys Lett A 1991; 155: 269–75.
- [4] Kudryashov N.A. Partial differential equations with solutions having movable first – order singularities, Phys Lett A 1992; 169: 237–42.
- [5] Vernov S.Yu. Constructing Solutions for the Generalized HenonHeiles System Through the Painleve Test, Theor Math Phys 2003; 135(3): 792–801.
- [6] Kudryashov N.A. Exact solitary waves of the Fisher equations, Phys Lett A 2005; 342: 99–106.
- [7] Hone A.N.W. Non–existence of elliptic travelling wave solutions of the complex Ginzburg–Landau equation, Physica D 2005; 205: 292–306.
- [8] Kudryashov N.A. Simplest equation method to look for exact solutions of nonlinear differential equations, Chaos Solitons and Fractals 2005; 24: 1217–31.
- [9] Vernov S.Yu. Proof of the Absence of Elliptic Solutions of the Cubic Complex Ginzburg–Landau Equation, Theor Math Phys 2006; 146(1): 131–139.
- [10] Kudryashov N.A., Demina M.V. Polygons of differential equations for finding exact solutions, Chaos Solitons and Fractals 2007; 33: 1480–96.
- [11] Kudryashov N.A., Loguinova N.B. Extended simplest equation method for nonlinear differential equations, Applied Mathematics and Computation 2008; 205: 396–402.
- [12] Kudryashov N.A. On ”new travelling wave solutions” of the KdV and the KdV–Burgers equations, Commun Nonlinear Sci Numer Simulat 2009; 14: 1891–900.
- [13] Kudryashov N.A., Loguinova N.B. Be careful with the Exp–function method, Commun Nonlinear Sci Numer Simulat 2009; 14: 1881–90.

- [14] Kudryashov N.A. Seven common errors in finding exact solutions of nonlinear differential equations, Commun Nonlinear Sci Numer Simulat 2009; 14: 3507–29.
- [15] Lavrent'ev M. A., Shabat B. V. Methods of theory of functions of complex variables. Moscow: Nauka, 1965. 716 p. (in Russian)
- [16] Eremenko A. Meromorphic traveling wave solutions of the Kuramoto-Sivashinsky equation, J Math Phys Anal Geom 2006; 2(3) 278–86.
- [17] Nevanlinna R. Eindeutige analytische Funktionen. Berlin: Springer, 1936. 353 p.
- [18] Sikorsky Yu.S. Elements of elliptic functions' theory with applications to mechanics. Moscow: Komkniga, 2006. 368 p. (in Russian)
- [19] Malfliet W., Hereman W. The Tanh method: I Exact solutions of non-linear evolution and wave equations, Phys Scripta 1996; 54: 563–8.
- [20] Parkes E.J., Duffy B.R. An automated tanh-function method for finding solitary wave solutions to non-linear evolution equations, Comput Phys Commun 1996; 98: 288–300.